

Map Unit Description

Rifle Area, Colorado, Parts of Garfield and Mesa Counties

50 Olney loam, 3 to 6 percent slopes

Setting

Elevation: 5000 to 6500 feet

Composition

Olney and similar soils: 85 percent

Description of Olney

Setting

Landform: Valley sides, alluvial fans

Down-slope shape: Convex, linear

Across-slope shape: Convex, linear

Parent material: Alluvium derived from sandstone and shale

Properties and Qualities

Slope: 3 to 6 percent

Drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate maximum: 15 percent

Gypsum maximum: 0 percent

Available water capacity: Moderate (about 7.6 inches)

Interpretive Groups

Land capability classification (irrigated): 3e

Land capability (non irrigated): 3c

Ecological site: Rolling Loam (R048AY298CO)

Typical Profile

0 to 12 inches: loam

12 to 33 inches: sandy clay loam

33 to 43 inches: gravelly sandy clay loam

43 to 60 inches: very gravelly sandy loam

Map Unit Description

Rifle Area, Colorado, Parts of Garfield and Mesa Counties

51 Olney loam, 6 to 12 percent slopes

Setting

Elevation: 5000 to 6500 feet

Composition

Olney and similar soils: 85 percent

Description of Olney

Setting

Landform: Valley sides, alluvial fans
Down-slope shape: Convex, linear
Across-slope shape: Convex, linear
Parent material: Alluvium derived from sandstone and shale

Properties and Qualities

Slope: 6 to 12 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 7.6 inches)

Interpretive Groups

Land capability classification (irrigated): 4e
Land capability (non irrigated): 4e
Ecological site: Rolling Loam (R048AY298CO)

Typical Profile

0 to 12 inches: loam
12 to 33 inches: sandy clay loam
33 to 43 inches: gravelly sandy clay loam
43 to 60 inches: very gravelly sandy loam

Map Unit Description

Rifle Area, Colorado, Parts of Garfield and Mesa Counties

56 Potts loam, 6 to 12 percent slopes

Setting

Elevation: 5000 to 7000 feet

Composition

Potts and similar soils: 85 percent

Description of Potts

Setting

Landform: Valley sides, benches, mesas

Down-slope shape: Convex, linear

Across-slope shape: Convex, linear

Parent material: Alluvium derived from basalt and/or alluvium derived from sandstone and shale

Properties and Qualities

Slope: 6 to 12 percent

Drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.60 in/hr)

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate maximum: 15 percent

Gypsum maximum: 0 percent

Available water capacity: High (about 10.3 inches)

Interpretive Groups

Land capability classification (irrigated): 4e

Land capability (non irrigated): 4e

Ecological site: Rolling Loam (R048AY298CO)

Typical Profile

0 to 4 inches: loam

4 to 28 inches: clay loam

28 to 60 inches: loam

Map Unit Description

Rifle Area, Colorado, Parts of Garfield and Mesa Counties

66 Torriorthents-Camborthids-Rock outcrop complex, steep

Setting

Landscape: Foothills
Elevation: 5000 to 8500 feet
Mean annual precipitation: 10 to 15 inches
Mean annual air temperature: 39 to 46 degrees F
Frost-free period: 80 to 105 days

Composition

Torriorthents, steep, and similar soils: 45 percent
Camborthids, steep, and similar soils: 20 percent
Rock outcrop, steep: 15 percent

Description of Torriorthents, steep

Setting

Landform: Mountainsides
Landform position (two-dimensional): Footslope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Stony, basaltic alluvium derived from sandstone and shale

Properties and Qualities

Slope: 15 to 70 percent
Depth to restrictive feature: 4 to 30 inches to Lithic bedrock
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 5 percent
Gypsum maximum: 0 percent
Available water capacity: Very low (about 2.4 inches)

Interpretive Groups

Land capability (non irrigated): 7e

Typical Profile

0 to 4 inches: variable
4 to 30 inches: fine sandy loam
30 to 34 inches: unweathered bedrock

Description of Camborthids, steep

Setting

Landform: Mountainsides
Landform position (two-dimensional): Footslope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Stony, basaltic alluvium derived from sandstone and shale

Properties and Qualities

Slope: 15 to 65 percent
Depth to restrictive feature: 15 to 60 inches to Lithic bedrock
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 10 percent
Gypsum maximum: 2 percent
Available water capacity: Low (about 4.0 inches)

Interpretive Groups

Land capability (non irrigated): 7e

Typical Profile

0 to 4 inches: variable