

# Map Unit Description

Weld County, Colorado, Southern Part

## 9 Ascalon loam, 1 to 3 percent slopes

### Setting

Elevation: 4500 to 4900 feet  
Mean annual precipitation: 12 to 17 inches  
Mean annual air temperature: 46 to 57 degrees F  
Frost-free period: 130 to 160 days

### Composition

Ascalon and similar soils: 85 percent  
Minor components: 15 percent

### Description of Ascalon

#### Setting

Landform: Terraces  
Down-slope shape: Linear  
Across-slope shape: Linear  
Parent material: Alluvium

#### Properties and Qualities

Slope: 1 to 3 percent  
Drainage class: Well drained  
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.20 to 2.00 in/hr)  
Frequency of flooding: None  
Frequency of ponding: None  
Calcium carbonate maximum: 10 percent  
Gypsum maximum: 0 percent  
Available water capacity: Moderate (about 8.0 inches)

#### Interpretive Groups

Land capability classification (irrigated): 2e  
Land capability (non irrigated): 4c

#### Typical Profile

0 to 8 inches: loam  
8 to 18 inches: sandy clay loam  
18 to 60 inches: fine sandy loam  
60 to 64 inches: fine sandy loam

### Minor Components

#### Platner

Percent of map unit: 7 percent

#### Olney

Percent of map unit: 5 percent

#### Stoneham

Percent of map unit: 3 percent

# Map Unit Description

Weld County, Colorado, Southern Part

## 46 Olney fine sandy loam, 0 to 1 percent slopes

### Setting

Elevation: 4600 to 5200 feet  
Mean annual precipitation: 11 to 15 inches  
Mean annual air temperature: 46 to 54 degrees F  
Frost-free period: 125 to 175 days

### Composition

Olney and similar soils: 85 percent  
Minor components: 15 percent

### Description of Olney

#### Setting

Landform: Plains  
Down-slope shape: Linear  
Across-slope shape: Linear  
Parent material: Mixed deposit outwash

#### Properties and Qualities

Slope: 0 to 1 percent  
Drainage class: Well drained  
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.57 to 2.00 in/hr)  
Frequency of flooding: None  
Frequency of ponding: None  
Calcium carbonate maximum: 15 percent  
Gypsum maximum: 0 percent  
Available water capacity: Moderate (about 7.0 inches)

#### Interpretive Groups

Land capability classification (irrigated): 3e  
Land capability (non irrigated): 4c  
Ecological site: Sandy Plains (R067BY024CO)

#### Typical Profile

0 to 10 inches: fine sandy loam  
10 to 20 inches: sandy clay loam  
20 to 25 inches: sandy clay loam  
25 to 60 inches: fine sandy loam

### Minor Components

#### Zigweid

Percent of map unit: 8 percent

#### Vona

Percent of map unit: 7 percent

# Map Unit Description

Weld County, Colorado, Southern Part

## 47 Olney fine sandy loam, 1 to 3 percent slopes

### Setting

Elevation: 4600 to 5200 feet  
Mean annual precipitation: 11 to 15 inches  
Mean annual air temperature: 46 to 54 degrees F  
Frost-free period: 125 to 175 days

### Composition

Olney and similar soils: 85 percent  
Minor components: 15 percent

### Description of Olney

#### Setting

Landform: Plains  
Down-slope shape: Linear  
Across-slope shape: Linear  
Parent material: Mixed deposit outwash

#### Properties and Qualities

Slope: 1 to 3 percent  
Drainage class: Well drained  
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.57 to 2.00 in/hr)  
Frequency of flooding: None  
Frequency of ponding: None  
Calcium carbonate maximum: 15 percent  
Gypsum maximum: 0 percent  
Available water capacity: Moderate (about 7.0 inches)

#### Interpretive Groups

Land capability classification (irrigated): 3e  
Land capability (non irrigated): 4c  
Ecological site: Sandy Plains (R067BY024CO)

#### Typical Profile

0 to 10 inches: fine sandy loam  
10 to 20 inches: sandy clay loam  
20 to 25 inches: sandy clay loam  
25 to 60 inches: fine sandy loam

### Minor Components

#### Zigweid

Percent of map unit: 10 percent

#### Vona

Percent of map unit: 5 percent