

Seyler B10-32D
Seyler B15-79HNM
Seyler B10-64-1HN

Map Unit Description

Weld County, Colorado, Southern Part

TSN- R64W - 6th pm
Sec. 10: NWSW
Weld County, CO

26 Haverson loam, 1 to 3 percent slopes

Setting

Elevation: 4500 to 4800 feet
Mean annual precipitation: 12 to 17 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 125 to 180 days

Composition

Haverson and similar soils: 85 percent
Minor components: 15 percent

Description of Haverson

Setting

Landform: Flood plains, stream terraces
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Stratified, calcareous alluvium

Properties and Qualities

Slope: 1 to 3 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.60 to 2.00 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 1 percent
Available water capacity: High (about 9.6 inches)

Interpretive Groups

Land capability classification (irrigated): 3e
Land capability (non irrigated): 4c
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 4 inches: loam
4 to 60 inches: stratified loamy sand to loam to clay loam

Minor Components

Vona

Percent of map unit: 5 percent

Heldt

Percent of map unit: 5 percent

Nunn

Percent of map unit: 5 percent

Seyler B10-32D
Seyler B15-79HNM
Seyler B10-64-1HN

Map Unit Description

Weld County, Colorado, Southern Part

TSN- R64W - 6th pm
Sec. 10: NWSW
Weld County, CO

51 Otero sandy loam, 1 to 3 percent slopes

Setting

Elevation: 4700 to 5250 feet
Mean annual precipitation: 12 to 15 inches
Mean annual air temperature: 48 to 52 degrees F
Frost-free period: 130 to 180 days

Composition

Otero and similar soils: 85 percent
Minor components: 15 percent

Description of Otero

Setting

Landform: Plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Eolian deposits and/or mixed outwash

Properties and Qualities

Slope: 1 to 3 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high or high (0.57 to 5.95 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 10 percent
Gypsum maximum: 0 percent
Available water capacity: Moderate (about 7.7 inches)

Interpretive Groups

Land capability classification (irrigated): 3e
Land capability (non irrigated): 4e
Ecological site: Sandy Plains (R067BY024CO)

Typical Profile

0 to 12 inches: sandy loam
12 to 60 inches: fine sandy loam

Minor Components

Kim

Percent of map unit: 10 percent

Vona

Percent of map unit: 5 percent

Map Unit Description

Weld County, Colorado, Southern Part

Seyler B10-32D
Seyler B15-79HNM
Seyler B10-64-1HN

TSN- R64W - 6th pm
Sec. 10: NWSW
Weld County, CO.

68 Ustic Torriorthents, moderately steep

Setting

Elevation: 4450 to 5100 feet
Mean annual precipitation: 10 to 16 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 120 to 160 days

Composition

Ustic torriorthents and similar soils: 85 percent
Minor components: 15 percent

Description of Ustic torriorthents

Setting

Landform: Escarpments, breaks
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Gravelly alluvium

Properties and Qualities

Slope: 9 to 15 percent
Drainage class: Excessively drained
Capacity of the most limiting layer to transmit water (Ksat): High or very high (5.95 to 19.98 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 5 percent
Gypsum maximum: 0 percent
Available water capacity: Very low (about 1.8 inches)

Interpretive Groups

Land capability (non irrigated): 7s

Typical Profile

0 to 10 inches: gravelly sand
10 to 60 inches: gravelly sand

Minor Components

Columbo

Percent of map unit: 10 percent

Eckley

Percent of map unit: 3 percent

Otero

Percent of map unit: 2 percent