

Map Unit Description

Weld County, Colorado, Northern Part

37 Midway clay loam, 0 to 9 percent slopes

Setting

Elevation: 3500 to 6000 feet
Mean annual precipitation: 10 to 17 inches
Mean annual air temperature: 46 to 52 degrees F
Frost-free period: 100 to 180 days

Composition

Midway and similar soils: 80 percent
Minor components: 20 percent

Description of Midway

Setting

Landform: Ridges, breaks, plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous clayey residuum weathered from shale

Properties and Qualities

Slope: 0 to 9 percent
Depth to restrictive feature: 6 to 20 inches to Paralithic bedrock
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 15 percent
Salinity maximum: Non saline or slightly saline (2.0 to 8.0 mmhos/cm)
Sodium adsorption ratio maximum: 15.0
Available water capacity: Very low (about 1.9 inches)

Interpretive Groups

Land capability classification (irrigated): 6e
Land capability (non irrigated): 6e
Ecological site: Shaly Plains (R067BY045CO)

Typical Profile

0 to 3 inches: clay loam
3 to 11 inches: clay
11 to 15 inches: weathered bedrock

Minor Components

Renohill

Percent of map unit: 9 percent

Stoneham

Percent of map unit: 6 percent

Terry

Percent of map unit: 5 percent

Rohn state LD09-65HN
Rohn state LD09-65-1HN
Rohn state LD09-64HN
Rohn state LD09-64-1HN
Rohn state LD10-65HN
Rohn state LD10-65-1HN
Rohn state LD10-64HN
Rohn state LD10-64-1HN

T9N-1258W, 6th p.m.
Sec. 9: NESE, SENE
Weld County, CO.

Map Unit Description

Weld County, Colorado, Northern Part

36 Manzanola clay loam, 0 to 3 percent slopes

Setting

Elevation: 4400 to 5600 feet
Mean annual precipitation: 11 to 15 inches
Mean annual air temperature: 46 to 52 degrees F
Frost-free period: 140 to 180 days

Composition

Manzanola and similar soils: 85 percent
Minor components: 15 percent

Description of Manzanola

Setting

Landform: Stream terraces, swales, plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous clayey alluvium

Properties and Qualities

Slope: 0 to 3 percent
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 5 percent
Gypsum maximum: 3 percent
Sodium adsorption ratio maximum: 15.0
Available water capacity: High (about 9.6 inches)

Interpretive Groups

Land capability (non irrigated): 4e
Ecological site: Clayey Plains (R067BY042CO)

Typical Profile

0 to 3 inches: clay loam
3 to 25 inches: clay
25 to 48 inches: clay
48 to 60 inches: clay loam

Minor Components

Avar

Percent of map unit: 15 percent

Rohn state LD09-65HN
Rohn state LD09-65HN
Rohn state LD09-64HN
Rohn state LD09-64-1HN
Rohn state LD10-65HN
Rohn state LD10-65-1HN
Rohn state LD10-64HN
Rohn state LD10-64-1HN

T9N-R58W, 6th PM
Sec. 9: NESE, SENE
Weld County, CO.

Map Unit Description

Weld County, Colorado, Northern Part

57 Renohill-Shingle complex, 3 to 9 percent slopes

Setting

Elevation: 3600 to 6200 feet
Mean annual precipitation: 10 to 16 inches
Mean annual air temperature: 46 to 48 degrees F
Frost-free period: 100 to 160 days

Composition

Renohill and similar soils: 50 percent
Shingle and similar soils: 35 percent
Minor components: 15 percent

Description of Renohill

Setting

Landform: Breaks, ridges, plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous, clayey loamy residuum weathered from shale

Properties and Qualities

Slope: 3 to 9 percent
Depth to restrictive feature: 20 to 40 inches to Paralithic bedrock
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: Low (about 5.0 inches)

Interpretive Groups

Land capability (non irrigated): 4e
Ecological site: Loamy Plains (R067BY002CO)

Typical Profile

0 to 4 inches: fine sandy loam
4 to 13 inches: clay
13 to 29 inches: clay loam
29 to 33 inches: unweathered bedrock

Description of Shingle

Setting

Landform: Breaks, plains, ridges
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous loamy residuum weathered from shale

Properties and Qualities

Slope: 3 to 9 percent
Depth to restrictive feature: 10 to 20 inches to Paralithic bedrock
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low or moderately high (0.06 to 0.20 in/hr)
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate maximum: 15 percent
Gypsum maximum: 0 percent
Available water capacity: Very low (about 2.1 inches)

Interpretive Groups

Land capability classification (irrigated): 6s
Land capability (non irrigated): 6s
Ecological site: Shaly Plains (R067BY045CO)

Typical Profile

Rohn state LD09-65HN
Rohn state LD09-65-1HN
Rohn state LD09-64HN
Rohn state LD09-64-1HN
Rohn state LD10-65HN
Rohn state LD10-65-1HN
Rohn state LD10-64HN
Rohn state LD10-64-1HN

T9N-R58W, 6th. pm
Sec 9: NESE, SENE
Weld County, CO.

Map Unit Description

Weld County, Colorado, Northern Part

0 to 4 inches: clay loam
4 to 11 inches: clay loam
11 to 15 inches: unweathered bedrock

Minor Components

Midway

Percent of map unit: 8 percent

Tassel

Percent of map unit: 7 percent