

Jackson County Area, Colorado

Sp—Spicerton sandy loam

Map Unit Setting

National map unit symbol: jq0p
Elevation: 7,800 to 8,300 feet
Mean annual precipitation: 10 to 12 inches
Mean annual air temperature: 36 to 38 degrees F
Frost-free period: 30 to 40 days
Farmland classification: Not prime farmland

Map Unit Composition

Spicerton and similar soils: 85 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Spicerton

Setting

Landform: Drainageways, stream terraces
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Coalmont formation alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 2 inches: sandy loam
H2 - 2 to 10 inches: clay
H3 - 10 to 60 inches: clay

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Moderately well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat):
Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Calcium carbonate, maximum in profile: 15 percent
Salinity, maximum in profile: Moderately saline to strongly saline
(8.0 to 16.0 mmhos/cm)
Sodium adsorption ratio, maximum in profile: 50.0
Available water storage in profile: Moderate (about 6.5 inches)

Interpretive groups

Land capability classification (irrigated): 6s
Land capability classification (nonirrigated): 7s
Hydrologic Soil Group: C
Ecological site: Salt Flats (R048AY261CO)

Hydric soil rating: No

Data Source Information

Soil Survey Area: Jackson County Area, Colorado
Survey Area Data: Version 7, Sep 22, 2014

Sp—Spicerton sandy loam						
Spicerton	Salt Flats	900	700	500	Western wheatgrass	15
					Inland saltgrass	15
					Alkali bluegrass	10
					Nuttall's alkaligrass	10
					Bottlebrush squirreltail	10
					Big sagebrush	10
					Winterfat	5
					Indian ricegrass	5
					Greasewood	5
Mat saltbush	5					